



ABOUT OUR HOSPITAL

The Perth and Smiths Falls District Hospital is a fully accredited health care facility located on two sites serving a catchment population of approximately 60,000+ residents within the tri-county region of Lanark, Leeds and Grenville

The hospital has a long history in the provision of high quality health services to its community.



60 Cornelia St W
Smiths Falls, ON
K7A 2H9
(613) 283 2330

33 Drummond St W,
Perth, ON
K7H 2K1
(613) 276 1500

www.psfhdh.on.ca

FERRIC DERISOMALTOSE (MONOFERRIC ©)

PATIENT INFORMATION



Patients and Families
THE Heart of Care

WHAT IS MONOFERRIC AND WHAT IS IT USED FOR?

It is an iron replacement therapy used to treat anemia (low red blood cells) caused by low iron levels.

WHAT SHOULD I TELL MY DOCTOR BEFORE RECEIVING MONOFERRIC?

- Your medical conditions
- Any medications or supplements you take
- If you've had an allergic reaction to iv iron before
- Are pregnant or plan to become pregnant
- Are breastfeeding or plan to breastfeed

HOW IS MONOFERRIC GIVEN?

It is given as an infusion into a vein over a period of time. You will be watched closely during and after this process, which can take anywhere from 30 minutes to 1 hour, depending on your dose.

WHAT TO DO AFTER AN INFUSION?

- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you
- To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs
- Have blood work checked as you have been told by the doctor

WHAT ARE SOME SIDE EFFECTS ONE MAY EXPERIENCE AFTER RECEIVING MONOFERRIC?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away.

- upset stomach
- headache
- constipation
- nose or throat irritation
- nausea, or throwing up

WHAT ARE SOME SERIOUS BUT RARE SIDE EFFECTS I NEED TO WATCH FOR AND CALL MY DOCTOR IF THEY OCCUR?

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat
- Signs of high or low blood pressure like very bad headache or dizziness, passing out, changes in eyesight
- This drug may stain your skin brown near where it is given if it leaks from the vein. This may last for a long time. Tell your nurse if you have any leaking of fluid from the injection site.